

2017

A new species of *Madoniella* Pic (Coleoptera: Cleridae) from Chiapas, Mexico

Jacques Rifkind

California State Collection of Arthropods

Weston Opitz

Florida State Collection of Arthropods

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

Rifkind, Jacques and Opitz, Weston, "A new species of *Madoniella* Pic (Coleoptera: Cleridae) from Chiapas, Mexico" (2017). *Insecta Mundi*. 1036.

<http://digitalcommons.unl.edu/insectamundi/1036>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0531

A new species of *Madoniella* Pic (Coleoptera: Cleridae)
from Chiapas, Mexico

Jacques Rifkind
California State Collection of Arthropods
3294 Meadowview Road
Sacramento, California, 95832 U.S.A.

Weston Opitz
Florida State Collection of Arthropods
1911 SW 34th Street
Gainesville, Florida, 32614-7100, U.S.A.

Date of Issue: March 31, 2017

Jacques Rifkind and Weston Opitz

A new species of *Madoniella* Pic (Coleoptera: Cleridae) from Chiapas, Mexico

Insecta Mundi 0531: 1–4

ZooBank Registered: urn:lsid:zoobank.org:pub:208F3625-547F-435F-8C91-756A8459795C

Published in 2017 by

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: David Plotkin, e-mail: insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and Submission Requirements available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

A new species of *Madoniella* Pic (Coleoptera: Cleridae) from Chiapas, Mexico

Jacques Rifkind

California State Collection of Arthropods, 3294 Meadowview Road,
Sacramento, California, 95832 U.S.A.
clerid@aol.com

Weston Opitz

Florida State Collection of Arthropods, 1911 SW 34th Street,
Gainesville, Florida, 32614-7100, U.S.A.
opitz@kwu.edu

Abstract. Opitz (2011) revised the New World clerid genus *Madoniella* Pic, and recognized 74 species. We describe and illustrate *Madoniella fitzgeraldae*, new species (Coleoptera: Cleridae), recently discovered in Chiapas, México.

Key Words. Taxonomy, tropical deciduous forest, oak forest.

Introduction

In his recent revision of the genus *Madoniella* Pic, Opitz (2011) recognized 74 species (65 new) primarily on the basis of distinctive elytral markings (the “elytral insignia”) that appear to be remarkably consistent and species-specific, at least for subtropical and tropical members of the genus. Many of these species were described from a single specimen or a small series, and many were collected only recently. As a genus, *Madoniella* has adapted to a broad range of habitats (e.g. pine and oak forests, tropical deciduous forest, broadleaved hardwood forest, rainforest, and cloud forest) but its component species often appear to have limited distributions, which may reflect restricted ecological tolerances. This high degree of endemism, coupled with these beetles’ small size and cryptic coloration, suggest that many more species remain to be discovered. One such species, recently collected in southern Mexico, is described herein.

Material and Methods

The dry pointed holotype specimen was photographed through the eyepiece of a Zeiss stereo dissecting microscope using the camera in an Apple iPhone 7 Plus. The specimen’s abdomen was removed and its pygidium dissected following techniques described in Ekis (1977). After examination, the pygidium and aedeagus were placed on a drop of gelatin in a microvial, and this affixed to the specimen’s pin. The holotype male is deposited in the Florida State Collection of Arthropods, Gainesville Florida, U.S.A. (FSCA); the single paratype will be deposited in the junior author’s collection (JNRC = Collection of Jacques Rifkind, Valley Village, California, U.S.A)

Taxonomy

Madoniella fitzgeraldae Rifkind and Opitz, new species
(Fig. 1–3)

Diagnosis. The new species can be distinguished from congeners by its small size (less than 3 mm in length) and the unique shape of the posterior block of the elytral insignia (Fig. 1). It appears most similar to *M. disjuga* Opitz, from which it differs by having the posterocentral extension of the insignial block reduced to a small triangulate streak on each elytron (Fig. 1). In *M. disjuga*, the posterocentral extension of the insignial block is subtruncate, and only slightly inflected at the middle of the posterior margin (Fig. 239 in Opitz 2011). In addition, *M. disjuga* appears to be confined to the central Mexican

states of San Luis Potosí and Querétaro, whereas *M. fitzgeraldae* is known from Chiapas, in extreme southern Mexico. Opitz's (2011) key to the species of *Madoniella* is modified as follows to include the new species:

- 50(49'). Posterocentral extension of insignial block reduced to a small, triangulate streak on each elytron (Mexico: Chiapas) ***Madoniella fitzgeraldae* new species.**
 50'. Posterocentral extension of insignial block fully formed **51**

Description. Length 2.8 mm. Color: cranium and pronotal disc piceous; antennae testaceous basally, capitulum infusate from anterior half of antennomere 8 to apex of antennomere 10; mouthparts reddish testaceous; elytra brownish black (paler on apices and lateral margins posteriorly), with a pattern of testaceous markings (elytral insignia) as in Fig. 1; legs testaceous, femora brownish dorsally. Head: vertex moderately wide; surface densely, coarsely punctate and rugulose; front moderately densely clothed with short, adpressed silvery setae. Pronotum: broader than long (ratio of width to length 5:4), anterior margin broadly arcuate; lateral margin generally convex (slightly inflected anteriorly); posterior angles rounded; disc convex above (Fig. 2), without an anterior transverse depression; surface punctation as on head; integument moderately densely but inconspicuously set with fine, rather short, suberect and reclinate grayish setae. Elytra: elongate (ratio of length to maximum width 93:40); form oblong–subovoid / oblong–rectangulate; surface covered with large, subquadrate punctations, arranged serially in 10 rows; raised interstitial areas between punctures forming shallow longitudinal carinae laterally; epipleural margin minutely serrate along posterior 1/3 (Fig. 2); integument moderately densely but inconspicuously vested with testaceous and black, rather short subreclinate and erect setae, not formed into matted clumps. Legs: protibial anterior margin with 4 spines. Abdomen: Comprised of 6 visible sternites; 6th visible sternite partially beneath 5th; pygidium transverse-scutiform, well sclerotized; aedeagus (Fig. 3) with tegmen tubular, weakly sclerotized, phallobasic lobes short, not fimbriate, phallobasic rod well developed, linear, phallobasic apodeme well developed, flared laterally at basal limit, phallobasic struts contiguous with phallobasic apodeme, phallic plates slender; spicular fork well developed, spicular plates flared, intraspicular plate linear, spicular apodemes fused from middle to basal limit.

Variation. The female paratype is quite similar to the holotype, with the exception of the shape of visible abdominal sternite 6, which is triangulate, with its hind margin rather narrowly rounded, and visible abdominal tergite 6, which has the hind margin moderately broadly subtruncate.

Type material. Holotype, male. MEXICO: Chiapas, 3.7 km NE Cacahuanó, 988 m, N 17°01.238' W 093°07.990', vi–25–2016, on slash, oak / thorn, J. Rifkind, E. Martinez, colls. Holotype deposited in FSCA. Paratype: 1 (JNRC), same data as holotype.

Etymology. The specific name is a patronymic in honor of Ella Fitzgerald, the great American jazz singer, often called “The First Lady of Song.”

Distribution. Chiapas, Mexico, in the municipality of Chicoasén, north of the city of Tuxla Gutiérrez.

Natural history. The holotype male and paratype female were collected *in copulo* on a slash pile in an ecotone between tropical deciduous forest and oak forest, at an elevation of 988 m. Within the immediate vicinity, a recently felled and cut up oak tree (*Quercus* sp.), slash of various thicknesses on the ground, and a standing, partially cut and sapping leguminaceous tree, had attracted several other species of Cleridae, as well as Cerambycidae, Buprestidae, Cantharidae, foraging *Camponotus* Mayr ants and a host of parasitic Hymenoptera.

Acknowledgments

The first author would like to thank Eric Antonio Martinez, of Mexico Birding Tours, the co-collector of the type series, for his friendship and energetic assistance in the field. Jesús Romero Nápoles

(Instituto de Fitosanidad; Colegio de Postgraduados, Montecillo, Estado de México, Mexico) once again generously provided official permission to collect in Chiapas. We appreciate comments on the manuscript from John Leavengood, Jr. and Henry Hespenheide. Finally, we thank Patricia Gum for her graciously offered help in preparing the illustrations.

Literature Cited

- Ekis, G. 1977.** Classification, phylogeny, and natural history of the genus *Perilypus* (Coleoptera: Cleridae). *Smithsonian Contributions to Zoology* 227: 1–138.
- Opitz, W. 2011.** Classification, natural history, and evolution of Epiphloeinae (Coleoptera, Cleridae); Part X. The genus *Madoniella* Pic, 1935. *Entomologica Basiliensia et Collectionis Frey* 33: 133–248.

Received January 29, 2017; Accepted February 15, 2017.
Review Editor Adam Brunke.



Figures 1–3. *Madoniella fitzgeraldae* new species. 1) habitus; 2) lateral aspect; 3) aedeagus.